

SEMI-LOCAL INVARIANTS FOR NON RESONANT POISSON STRUCTURES ON $S^1 \times \mathbb{R}^n$.

OLIVIER BRAHIC JEAN-PAUL DUFOUR.

ABSTRACT. The local study of a Poisson structure near a “generic” isolated zero has been the subject of many works. The purpose of this paper is to give a “semi-local” study along “generic” closed curves of such zeros: we formally classify Poisson structures defined in a neighborhood of $\Gamma = S^1 \times \{0\}$ in $S^1 \times \mathbb{R}^n$, that vanish on Γ , and whose linear approximation at one point m of Γ is isomorphic to the dual of a non-resonant Lie algebra.

1. INTRODUCTION.

Let Π be a Poisson structure on some manifold M , vanishing at a point $m \in M$. Denote $\mathcal{D}_\omega \Pi$ the modular vector field ([W2]) with respect to some density ω , and suppose that $\mathcal{D}_\omega \Pi$ doesn't vanish at m ; this requirement is independent of the chosen density, and forces the Poisson structure to vanish all along the orbit Γ of $\mathcal{D}_\omega \Pi$ through m , thus Γ is a whole curve of singularities for Π . In this paper, we are interested in describing generic Poisson structure in the neighborhood of such Γ , when it is supposed to be closed, that is $\Gamma = S^1$. Also we will consider that Π is defined on a neighborhood of $\Gamma = S^1 \times \{0\}$ in $S^1 \times \mathbb{R}^n$ parametrized by $(\theta, x_1, \dots, x_n)$, and that Π is zero on Γ .

The first invariant attached to this situation is the period c of this orbit Γ : it doesn't depend on the chosen density because Π vanishes on it. Recall that this is one of the invariants used by O. Radko ([R]) to classify Poisson structures on surfaces.

The second thing we have to do is to take care of the linear part of Π at the points of Γ . If we choose local coordinates $(\tilde{x}_0, \tilde{x}_1, \dots, \tilde{x}_n)$ vanishing at m , and such that $\mathcal{D}_\omega \Pi$ is $\frac{\partial}{\partial \tilde{x}_0}$, this linearized Poisson structure satisfies relations:

$$\begin{aligned} \{\tilde{x}_0, \tilde{x}_i\} &= \sum_{j=1}^n a_{i,j} \tilde{x}_j \\ \{\tilde{x}_r, \tilde{x}_i\} &= \sum_{j=1}^n c_{r,i}^j \tilde{x}_j, \end{aligned}$$

for r and i varying from 1 to n . This means that it corresponds to a Lie algebra (see [W1]) which is a semi-direct product of $\mathbb{R}$ with a n -dimensional Lie algebra A . In the sequel we will denote by D the derivation of A under which $\mathbb{R}$ acts on A : in the above coordinates, it has the matrix with coefficients $a_{i,j}$. We see also that, up to a linear isomorphism, this Lie algebra doesn't depend on the point m chosen on Γ as $\mathcal{D}_\omega \Pi$ is an infinitesimal isomorphism. So this isomorphism class of Lie algebra is the second invariant we can attach to our situation.

The generic condition we will impose on Π in the sequel, is that there are no resonance relation between the eigenvalues $\lambda_1, \dots, \lambda_n$ of D ; by this we mean precisely that there are no relations

$$\lambda_i = p_1 \lambda_1 + \dots + p_n \lambda_n,$$

$$\lambda_i + \lambda_j = p_1 \lambda_1 + \cdots + p_n \lambda_n$$

for every i and j , ($i \neq j$), where $(p_1, \dots, p_n)$ is a multi-index with non-negatives p_k and $\sum_{k=1}^n p_k \geq 2$, except trivial relations $\lambda_i + \lambda_j = \lambda_i + \lambda_j$. In particular, these eigenvalues are *distinct*, and D is diagonalizable. It is then easy to verify that, in the complex case, the brackets of such linear Poisson structures, so called **non resonant**, and denoted $(\mathbb{C}^{n+1})_{(\lambda_1 \dots \lambda_n)}^*$, can always be written as:

$$\begin{aligned} \{e_i, e_j\} &= 0, & i, j &= 1 \dots n \\ \{e_0, e_i\} &= \lambda_i e_i, & i &= 1 \dots n, \end{aligned}$$

(the derivation D is just $\{e_0, -\}$) so that A has to be commutative.

Note that two such Lie algebras $(\mathbb{C}^{n+1})_{(\lambda_1 \dots \lambda_n)}^*$ and $(\mathbb{C}^{n+1})_{(\lambda'_1 \dots \lambda'_n)}^*$ are isomorphic if and only if $\lambda_i = d \lambda'_i$, up to a re-indexation of the λ_i 's, and for $d \in \mathbb{C}$ (see the proof of lemma 2.1). Also, in terms of classification of Lie algebras, the λ_i 's should be seen as a set defined up to multiplication by a scalar of all its elements.

In [D-Z] there is a local study of this situation (in the real and complex cases): it appears there that the Poisson structure is, at least formally, "quadratzable", i.e. we can find coordinates such that our Poisson structure has only linear and quadratic terms. In this paper we will show that this is also true in a neighborhood of the whole curve Γ . Moreover we will describe all the invariants attached to the isomorphism class of the germ of Π along Γ . However, in order to simplify the study, we will restrict ourselves to the case where **the eigenvalues λ_i are all real**. In that case the Lie algebra has the above form but in real coordinates; it will be then denoted by $(\mathbb{R}^{n+1})_{(\lambda'_1 \dots \lambda'_n)}^*$.

In the analytic context, we will need an extra assumption on these eigenvalues : we will suppose that they verify Bruno's condition, that is : if I denote the set of all mutli-indexes $(c_i)_{i=1 \dots n} \in \mathbb{Z}^n$ such that $c_i \leq -1, \forall i = 1 \dots n$ and $\sum_{i=1 \dots n} c_i \lambda_i \neq 0$, one puts :

$$(1.1) \quad \omega_k = \min\{|\sum_{i=1}^n c_i \lambda_i|; (c_i) \in I, \sum_{i=1}^n c_i < 2^k\},$$

Imposong Bruno's condition is requiring that $\sum_{k=1}^{\infty} \frac{1}{2} \log \frac{1}{\omega_k} < \infty$, it is a standard condition for assuring analytic convergence of formal series.

It is willingly that we didn't precise if Π was supposed to be smooth or analytic, next section is valid for the smooth case, and also, under Bruno's condition, in the analytic case (of course, flat terms $o_{\infty}(x)$ will have to be omitted in the analytic case).

2. THE NORMAL FORM.

In this section, we give several lemmas, that will, step by step, lead to a normal form for Π in a neighborhood of Γ , with underlying preoccupation to obtain coordinates that are global with respect to $\theta \in S^1$.

Lemma 2.1. *The brackets induced by Π , can, in a neighborhood of Γ , be written as:*

$$\begin{aligned} \{x_i, x_j\} &= o_2(x), & i, j = 1 \dots n \\ \{\theta, x_i\} &= \sum_{j=1}^n h_{i,j} x_j + o_2(x) & i = 1 \dots n, \end{aligned}$$

where $h_{i,j}$ are smooth functions on S^1 , $o_2(x)$ denote smooth functions on $S^1 \times \mathbf{R}^n$ and of order two in the variables $x_1, \dots, x_n$, and the matrix $H_\theta = (h_{i,j}(\theta))_{i,j=1 \dots n}$ has eigenvalues $\{k(\theta)\lambda_i/i = 1 \dots n\}$ for some $k : S^1 \mapsto \mathbf{R}$.

Proof. Π vanishes on $\Gamma = \{x_i = 0 ; i = 1 \dots n\}$, so that we can write the brackets as:

$$\begin{aligned} \{x_i, x_j\} &= \sum_{k=1 \dots n} u_k^{i,j}(\theta) x_k + o_2(x), & i, j = 1 \dots n \\ \{\theta, x_i\} &= \sum_{j=1 \dots n} h_{i,j}(\theta) x_j + o_2(x), & i = 1 \dots n. \end{aligned}$$

In fact the lemma is just a consequence of the fact that two semi-direct products of Lie algebras $\mathbf{R}_L \ltimes \mathbf{R}^n$ and $\mathbf{R}_{L'} \ltimes \mathbf{R}^n$ are isomorphic if and only if L and L' are conjugated up to a scalar, but let's see this in detail.

Let $\theta_0 \in S^1$ fixed, denote $\Pi_{\theta_0}^{(1)}$ the linear approximation of Π at $(\theta_0, 0, \dots, 0)$, according to the introduction, it is isomorphic to the dual of a Lie algebra with commutative derived ideal, so all $u_k^{i,j}(\theta)$ must be zero.

Now, let $\Psi_{\theta_0} : (\mathbf{R}^3)_{(\lambda_1 \dots \lambda_n)}^* \mapsto \Pi_{\theta_0}^{(1)}$ a Poisson isomorphism, and

$$\tilde{\Psi}_{\theta_0} = \begin{pmatrix} k & v_1 & \dots & v_n \\ w_1 & & & \\ \vdots & & \Psi_{i,j} & \\ w_n & & & \end{pmatrix}$$

the associated matrix; $(\Psi_{\theta_0})^*$ being a Lie algebra isomorphism, it respects derived ideals, and $(w_1, \dots, w_n)$ must be zero.

Then we see that $\det(\Psi_{\theta_0}) = k \cdot \det(\psi_{i,j})_{i,j}$, in particular $k \neq 0$ and $(\Psi_{i,j})_{i,j}$ is invertible.

At that point, the conditions for Ψ_{θ_0} to be Poisson are:

$$\sum_{k=1, \dots, n} h_{i,k} \psi_{k,j} = k \cdot \psi_{i,j} \lambda_i, \quad i, j = 1 \dots n,$$

and we see that these relations can be written as:

$$H_{\theta_0} \cdot (\psi_{i,j}) = (\psi_{i,j}) \cdot \begin{pmatrix} k \cdot \lambda_n & & \\ & \ddots & \\ & & k \cdot \lambda_n \end{pmatrix}.$$

So the result. $\square$

Lemma 2.2. *Up to a covering, and in a well-fitted coordinate chart, the brackets induced by Π can be written in a neighborhood of Γ as:*

$$\begin{aligned} \{x_i, x_j\} &= o_2(x) \\ \{\theta, x_i\} &= k(\theta) \lambda_i + o_2(x) \quad \forall i = 1 \dots n, \end{aligned}$$

where $k \in C^\infty(S^1)$.

Proof. For any $\theta \in S^1$, H_θ has distinct eigenvalues $\{k(\theta)\lambda_i ; i = 1 \dots n\}$ so k is smooth, and, if we denote E_{λ_i} the associated characteristic spaces, they are supplementary subvectorbundles of $S^1 \times \mathbb{R}^n \mapsto S^1$, so that each of them is either trivial, or diffeomorphic to the Möbius band M .

The covering

$$\begin{aligned} C : \mathbf{S}^1 \times \mathbb{R}^n &\mapsto \mathbf{S}^1 \times \mathbb{R}^n \\ (\tilde{\theta}, \tilde{x}) &\mapsto (2\tilde{\theta}, \tilde{x}), \end{aligned}$$

induces on the source space a Poisson structure $\tilde{\Pi}$ for which the associated characteristic bundles $E_{\tilde{\lambda}_i}$ are all trivial. As the topology of these subbundles is invariant by Poisson isomorphisms, it is clear that two germs of Poisson structures Π and Π' on $S^1 \times \mathbb{R}^n$ are isomorphic if and only each E_{λ_i} is diffeomorphic to one of the $E_{\lambda'_i}$'s and $\tilde{\Pi}$ is Poisson isomorphic to $\tilde{\Pi}'$.

Also, up to a covering, E_{λ_i} is trivial for any i . Then the choice of non vanishing smooth sections for each of them defines a linear change of coordinates that diagonalizes the matrix H_θ . $\square$

Remark 2.3. Of course, Poisson structures on $S^1 \times \mathbb{R}^n$ admitting non trivial characteristic subbundles do exist, the basic exemple on $S^1 \times \mathbb{R}^2$ is given by the following brackets:

$$\begin{aligned} \{x_1, x_2\} &= \frac{x_1^2}{2}(\lambda \cos^2 \frac{\theta}{2} + \mu \sin^2 \frac{\theta}{2}) + \frac{x_2^2}{2}(\lambda \sin^2 \frac{\theta}{2} + \mu \cos^2 \frac{\theta}{2}) + x_1 x_2(\mu - \lambda) \cos \frac{\theta}{2} \sin \frac{\theta}{2} \\ \{\theta, x_1\} &= x_1(\lambda \cos^2 \frac{\theta}{2} + \mu \sin^2 \frac{\theta}{2}) + x_2(\mu - \lambda) \cos \frac{\theta}{2} \sin \frac{\theta}{2} \\ \{\theta, x_2\} &= x_1(\mu - \lambda) \cos \frac{\theta}{2} \sin \frac{\theta}{2} + x_2(\lambda \sin^2 \frac{\theta}{2} + \mu \cos^2 \frac{\theta}{2}). \end{aligned}$$

Here, it is interesting to notice how these brackets simplify when $\lambda = \mu = 1$.

Lemma 2.4. *By re-parametrizing S^1 , one can put the brackets under the following form:*

$$\begin{aligned} \{x_i, x_j\} &= o_2(x) \\ \{\theta, x_i\} &= \mu_i x_i + o_2(x), \quad i = 1 \dots n, \end{aligned}$$

Proof. The formula

$$\chi(\theta) = \frac{2\pi}{\int_0^{2\pi} k(t) dt} \int_0^\theta k^{-1}(t) dt$$

defines a diffeomorphism $S^1 \mapsto S^1$ that gives the announced re-parametrization. $\square$

Lemma 2.5. *There exists a coordinate chart, defined on a neighborhood of Γ , in which the brackets express as:*

$$\begin{aligned} \{x_i, x_j\} &= o_2(x) \\ \{\theta, x_i\} &= \mu_i x_i, \quad i = 1 \dots n. \end{aligned}$$

Proof. Let X_θ the hamiltonian vector field associated to the (multi-)function θ :

$$X_\theta = \sum_{i=1 \dots n} (\mu_i x_i + p_i) \frac{\partial}{\partial x_i},$$

where p_i are $o_2(x)$. This vector field turns out to be linearisable, considered as a vector field in the variables x with parameter θ : there is a family of local diffeomorphisms ϕ_θ , depending smoothly on θ , such that

$$\phi_{\theta*} X_\theta = \sum_{i=1 \dots n} \mu_i x_i \frac{\partial}{\partial x_i}$$

for every θ . Moreover we can choose ϕ_θ 1-periodic in θ . So the change of coordinates $(x, \theta) \mapsto (\phi_\theta(x), \theta)$ gives the result. $\square$

Remark 2.6. Let us give an idea of how to prove that X_θ is linearisable: first notice that a diffeomorphism $\phi_\theta = (\phi_\theta^1, \dots, \phi_\theta^n)$ linearises X_θ if and only if

$$\mathcal{L}_{X_\theta} \phi_\theta^i = \mu_i \phi_\theta^i \quad \forall i = 1 \dots n, \forall \theta \in S^1.$$

The first step consists in proving the result formally: write X_θ as

$$X_\theta = X_\theta^{(1)} + X_\theta^{(r)} + o_r(\mathbf{x}),$$

with $X_\theta^{(1)}$, $X_\theta^{(r)}$ and $o_r(x)$ denoting respectively the terms of order 1, of order $r > 1$ and of order higher than r in the x_i 's variables, and show the existence of a formal diffeomorphism $\phi_\theta = Id + \sum_{i \in \mathbb{N}} \phi_\theta^{(r)}$ by induction on r , using non-resonance conditions. Then, under Bruno's condition, one can show that the series ϕ_θ converge, getting the result in the analytic context. In the smooth case, one can proceed as in [RS]: what precedes, leads us to the form $X_\theta = X_\theta^{(1)} + o_\infty(x)$ where $o_\infty(x)$ denotes a vector field flat at $0 \in \mathbb{R}^n$ for all $\theta \in S^1$, it is then possible to eliminate these flat terms by direct integration, flatness assuring the convergence of this integral.

Lemma 2.7. *A last change of coordinates leads to the following brackets:*

$$\begin{aligned} \{x_i, x_j\} &= a_{i,j} x_i x_j + o_\infty(x), & i, j &= 1 \dots n \\ \{\theta, x_i\} &= \mu_i x_i, & i &= 1 \dots n. \end{aligned}$$

Proof. Let $u_{i,j} = \{x_i, x_j\}$, Jacobi's identity $\oint_{\theta, x_i, x_j} \{\theta, \{x_i, x_j\}\} = 0$ is now just:

$$(\mu_i + \mu_j) u_{i,j} = X_\theta(u_{i,j}).$$

First notice that $u_{i,j}^0 = x_i x_j$ is a particular solution of this equation, and that $u_{i,j}$ being another solution, writing it as $u_{i,j} = w_{i,j} x_i x_j$, $w_{i,j}$ has to be a first integral of X_θ .

Now, because of the dynamic of this vector field, the restriction $(w_{i,j})|_{\bigcup E_{\mu_i}}$ only depends on θ so $w_{i,j} = k_{i,j}(\theta) + \tau_{i,j}$, where $\tau_{i,j}$ is flat on Γ (notice that if all the μ_i 's have the same sign, then $\tau_{i,j}$ is necessarily zero because it is a first integral of X_θ).

Writing the terms of order one in Jacobi's identity $\oint_{x_1, x_i, x_j} \{x_1, \{x_i, x_j\}\} = 0$ with $k_{1,i}$ and $k_{1,j}$ independent of θ implies that $k_{i,j}$ is independent of θ too, so we only have to find a change of coordinates making $k_{1,i}$ independent of θ for any $i = 1 \dots n$. This diffeomorphism can be defined by:

$$(\theta, x_1, \dots, x_n) \mapsto (\theta, x_1, \chi_2 x_2, \dots, \chi_n x_n),$$

where

$$\chi_j(\theta) = e^{\frac{1}{\lambda_1} \int_0^\theta (k_{1,j}(t) - \bar{k}_{1,j}) dt}$$

with $\bar{k}_{1,j} = \frac{1}{2\pi} \int_0^{2\pi} k_{1,j}(t) dt$ for any $j = 2 \dots n$. $\square$

Remark 2.8. In the smooth context, when $n = 2$, it is still possible to improve this result by eliminating flat terms, the proof is rather technical and specific to the case $n = 2$, it can be found in [Br].

Remark 2.9. In the case $n = 1$ the normal form reduces to

$$\pi = cx \frac{\partial}{\partial \theta} \wedge \frac{\partial}{\partial x}$$

and we obtain, as a particular case, the form given in [R].

In the case $n = 2$ we can obtain a slightly better result: the (analytic and smooth) normal form is

$$\pi = \frac{\partial}{\partial \theta} \wedge (\mu_1 x_1 \frac{\partial}{\partial x_1} + \mu_2 x_2 \frac{\partial}{\partial x_2}) + ax_1 x_2 \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial x_2}$$

where a , and μ_1, μ_2 are real constants.

3. GEOMETRIC SIGNIFICANCE OF THE PARAMETERS.

In that section we will see that the parameters μ_i and $a_{i,j}$, which appear in the above normal form, are invariants of the Poisson structure (by this we mean that two equivalent such Poisson structures must have normal forms with the same μ_i 's and $a_{i,j}$'s) and give them a geometrical interpretation. The sum of the μ_i 's gives the period c of the modular vector field along the singular curve, the n -tuple $(\mu_1, \dots, \mu_n)$ characterises, up to a multiplicative constant, the linear approximation of Π along Γ , finally the μ_i 's and $a_{i,j}$'s measure the (non Poisson) holonomy of the symplectic foliation when we move along this singular curve.

- *Period of the rotational and linear approximation along Γ*

The modular vector field (see [W1]) $D_\omega \Pi$ of Π with respect to the volume form $\omega = d\theta \wedge dx_1 \wedge \dots \wedge dx_n$ (coordinates of the normal form) has the following expression:

$$D_\omega \Pi = \left(\sum_{i=1 \dots n} \mu_i + o_1(x) \right) \frac{\partial}{\partial \theta} + o_2(x).$$

As $D_\omega \Pi$ is defined up to hamiltonian vector fields, its restriction to Γ depends only on Π , but $2\pi / \sum_{i=1 \dots n} \mu_i$ appears to be the period of its flow on Γ . So, as we have remarked in the introduction, it is an invariant of the Poisson structure.

We have seen that, at a purely linear level, the set $\{\mu_1, \dots, \mu_n\}$ was defined up to a multiplicative constant. Because the linear part of the Poisson structure along the singular curve is (up isomorphism) an invariant this set, up to multiplication by a constant, is also an invariant. If we use the preceding result we obtain that the μ_i themselves are invariants. We recall that, in the local study ([DZ]) the μ_i are only invariant up to multiplication by a constant. Also this local study shows that the μ_i (up to multiplication by a constant) measure the way the symplectic leaves behave on a transversal to the singular curve: the traces of these leaves on such a transversal give a figure diffeomorphic to the phase portrait of the linear vector field $\sum_{i=1 \dots n} \mu_i x_i \partial / \partial x_i$.

- *The symplectic foliation*

Before showing how the $a_{i,j}$'s are involved in the symplectic foliation of Π , let us show that they are Poisson invariants : consider two Poisson structures Π and Π' on a neighborhood of Γ , and Φ a Poisson isomorphism from Π to Π' . Put Π and Π' under normal form, as the μ_i 's are Poisson invariants, we can suppose that

$\mu_i = \mu'_i$ for any $i = 1 \dots n$. Then, writing down the equations for Φ to be Poisson, we get, with evident notations, $\forall i, j = 1 \dots n$:

$$\begin{aligned} \sum_{k=1}^n \mu_k x_k \left(\frac{\partial \Phi_0}{\partial \theta} \frac{\partial \Phi_i}{\partial x_k} - \frac{\partial \Phi_0}{\partial x_k} \frac{\partial \Phi_i}{\partial \theta} \right) + \sum_{k,l=1}^n a_{k,l} x_k x_l \left(\frac{\partial \Phi_0}{\partial x_k} \frac{\partial \Phi_i}{\partial x_l} \right) &= \mu_i \Phi_i \quad (I)_i \\ \sum_{k=1}^n \mu_k x_k \left(\frac{\partial \Phi_i}{\partial \theta} \frac{\partial \Phi_j}{\partial x_k} - \frac{\partial \Phi_i}{\partial x_k} \frac{\partial \Phi_j}{\partial \theta} \right) + \sum_{k,l=1}^n a_{k,l} x_k x_l \frac{\partial \Phi_i}{\partial x_k} \frac{\partial \Phi_j}{\partial x_l} &= a'_{i,j} \Phi_i \Phi_j \quad (II)_{i,j}. \end{aligned}$$

Differentiating equation $(I)_i$ with respect to θ , and evaluating it onto Γ , we obtain:

$$\frac{\partial \Phi_i}{\partial \theta}(\theta, 0, 0) = 0 \quad \forall \theta \in S^1.$$

Differentiating it with respect to x_j , and evaluating it onto Γ , we get:

$$\mu_j \left(\frac{\partial \Phi_0}{\partial \theta} \frac{\partial \Phi_i}{\partial x_j} - \frac{\partial \Phi_0}{\partial x_j} \frac{\partial \Phi_i}{\partial \theta} \right)(\theta, 0, 0) = \mu_i \frac{\partial \Phi_i}{\partial x_j}(\theta, 0, 0) \quad \forall \theta \in S^1.$$

But $\Phi|_\Gamma$ must exchange modular vector fields, so $\Phi_1(\theta, 0, 0) = \theta + \text{constant}$ and this equation becomes:

$$\frac{\partial \Phi_i}{\partial x_j}(\theta, 0, 0) = 0 \quad \forall \theta \in S^1.$$

In conclusion, we can write Φ as:

$$\begin{aligned} \Phi_0 &= \theta + o_2(x) \\ \Phi_i &= u_i(\theta) x_i + o_2(x) \quad \forall i = 1 \dots n \end{aligned}$$

And in $(II)_{i,j}$, the terms of order two in the x_i 's variables give the relation:

$$\frac{\partial}{\partial \theta}(\mu_j \ln |u_i| + \mu_i \ln |u_j|) = a_{i,j} - a'_{i,j}$$

Here, the left side is the derivative of a function defined on S^1 , so its integral over S^1 is zero, and $a_{i,j} = a'_{i,j}$.

Let us now describe the symplectic foliation of Π , from on (in the smooth case), it will be supposed that *no flat terms appear in the normal form*, in other words, we suppose that

$$\Pi = \partial \theta \wedge \left(\sum_{i=1 \dots n} \mu_i x_i \partial x_i \right) + \sum_{1 \leq i < j \leq n} a_{i,j} x_i x_j \partial x_i \wedge \partial x_j.$$

We will only describe the symplectic foliation on the open set $P^+ := \{(\theta, x) \in S^1 \times \mathbb{R}^n / x_i > 0, \forall i = 1 \dots n\}$, however, the following reasoning allows to describe it entirely: indeed if, for any subset $I \subset \{1, \dots, n\}$ we denote $P_I = \{(\theta, x) \in S^1 \times \mathbb{R}^n / x_i = 0, \forall i \notin I\}$, one easily sees that P_I is a Poisson submanifold of the presently studied type (that contains itself Poisson submanifolds P_J with $J \subset I$), so we can apply what follows to $P_I^+ := \{(\theta, x) \in S^1 \times \mathbb{R}^n / x_i = 0, \forall i \notin I, x_i > 0 \text{ else}\}$. Moreover, it will appear clearly that P^+ is one of the connected component of the regular open set of Π —all these component being isomorphic as the reflexions $(\theta, x_1, \dots, x_i, \dots, x_n) \mapsto (\theta, x_1, \dots, -x_i, \dots, x_n)$ are Poisson—and that the complement of this regular set is precisely composed by the P_I for $I \subset \{1 \dots n\}, I \neq \{1 \dots n\}$.

To put the foliation induced by Π on P^+ in evidence, one first apply the diffeomorphism

$$\begin{aligned} L : \quad P^+ &\longmapsto S^1 \times \mathbb{R}^n \\ (\theta, x_1, \dots, x_n) &\longmapsto (\theta, \ln x_1, \dots, \ln x_n), \end{aligned}$$

as

$$\{\theta, \ln x_i\} = \mu_i \{\ln x_i, \ln x_j\} = a_{i,j},$$

in the induced coordinates (that we shall denote by $(\theta, \bar{\mathbf{x}}) = (\theta, \bar{x}_1, \dots, \bar{x}_n) = (\theta, \ln x_1, \dots, \ln x_n)$), the matrix $(\Pi_{i,j})$ of the brackets has constant coefficients:

$$(\Pi_{i,j}) = \begin{pmatrix} 0 & -\mu_1 & \dots & -\mu_n \\ \mu_1 & & & \\ \vdots & & (a_{i,j}) & \\ \mu_n & & & \end{pmatrix}.$$

Denote μ the vector $(\mu_1, \dots, \mu_n)$ and $2s$ the rank of the matrix $(a_{i,j})$, the theorem of spectral decomposition for anti-autoadjoint operators assures the existence of a linear isomorphism $\phi : \mathbb{R}^n \mapsto \mathbb{R}^n$, whose matrix will be denoted $(\phi_{i,j})$ that conjugates $(a_{i,j})$ to a matrix of the following type

$$\begin{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} & & O & O \\ & \ddots & & \vdots \\ O & & \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} & O \\ O & \dots & O & \begin{pmatrix} 0 & \dots & 0 \\ \vdots & & \vdots \\ 0 & \dots & 0 \end{pmatrix} \end{pmatrix}.$$

Moreover, if μ belongs to $\text{Im}(a_{i,j})$, one can choose μ as an element of the basis for which $(a_{i,j})$ takes that form, this incites us to distinguish two cases :

1st case: if $\mu \in \text{Im}(a_{i,j})$

Choosing ϕ as described below, the diffeomorphism $(\theta, \mathbf{x}) \mapsto (\theta, \phi(\mathbf{x}))$ conjugates $(\Pi_{i,j})$ to the following matrix:

$$\begin{pmatrix} 0 & 0 & -1 & & \dots & & 0 \\ 0 & \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} & & & O & & \\ 1 & & & & & & \\ \vdots & & \ddots & & & & \vdots \\ \vdots & O & & \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} & & & \\ 0 & & \dots & 0 & & \begin{pmatrix} 0 & \dots & 0 \\ \vdots & & \vdots \\ 0 & \dots & 0 \end{pmatrix} \end{pmatrix}.$$

Here, the null matrix, in the lower right corner is of order $n - 2s$.

Denote $(\theta, q_1, p_1, \dots, q_s, p_s, \tilde{x}_{2s+1}, \dots, \tilde{x}_n)$ the coordinates induced by $\phi \circ L$, the foliation is then spanned by $\partial\theta + \partial q_1, \partial p_1, \partial q_2, \partial p_2, \dots, \partial q_s, \partial p_s$, so that this foliation is not parallel to $S^1 \times \{0\}$, letting a holonomy phenomenon appear along Γ . The set of leaves can be parametrised by $(q_1, \tilde{x}_{2s+1}, \dots, \tilde{x}_n) \in [0, 2\pi[\times \mathbb{R}^{n-2s}$.

Moreover, if $\mathcal{F}_{(q_1, \tilde{x}_{2s+1}, \dots, \tilde{x}_n)}$ denotes the leaf through the point $(0, q_1, 0, \dots, 0, \tilde{x}_{2s+1}, \dots, \tilde{x}_n) \in S^1 \times \mathbb{R}^n$, then $\mathcal{F}_{(q_1, \tilde{x}_{2s+1}, \dots, \tilde{x}_n)}$ is parametrised by $(t_1, \dots, t_{2s})$:

$$\mathcal{F}_{(q_1, \tilde{x}_{2s+1}, \dots, \tilde{x}_n)} = \{(t_1, q_1 + t_1, t_2, t_3, \dots, t_{2s}, \tilde{x}_{2s+1}, \dots, \tilde{x}_n) \in S^1 \times \mathbb{R}^n / t_1 \dots t_{2s} \in \mathbb{R}\},$$

In the coordinates of the normal form, it appears that the leaf $F_{(x_1, \dots, x_n)}$ through the point $(0, x_1, \dots, x_n)$ admits the following parametrisation:

$$F_{(x_1, \dots, x_n)} = \{(t_1, (x_i e^{\sum_{j=1 \dots 2s} \psi_{i,j} t_j})_{i=1 \dots n}) \in P^+ / t_1 \dots t_{2s} \in \mathbb{R}\},$$

where $(\psi_{i,j})$ denotes the inverse matrix of $(\phi_{i,j})$.

2nd case: if $\mu \notin \text{Im}(a_{i,j})$

We follow the same reasoning: L défnit a chart on P^+ in which the matrix $(\Pi_{i,j})$ can be conjugated by a linear isomorphism ϕ to the following

$$\begin{pmatrix} 0 & 0 & 0 & \dots & -1 & \dots \\ \vdots & \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} & & O & & \\ & & \ddots & & & \vdots \\ \vdots & O & & \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} & & \\ 1 & & & & \begin{pmatrix} 0 & \dots & 0 \\ \vdots & & \vdots \\ 0 & \dots & 0 \end{pmatrix} \\ \vdots & & & & & \\ 0 & & \dots & 0 & & \end{pmatrix}.$$

In the coordinates induced by $\phi \circ L$, that we shall denote by $(\theta, q_1, p_1, \dots, q_s, p_s, \tilde{x}_{2s+1}, \dots, \tilde{x}_n)$, the foliation is spanned by $\partial\theta, \partial q_1, \partial p_1, \partial q_2, \partial p_2, \dots, \partial q_s, \partial p_s, \partial \tilde{x}_{2s+1}$, so it is parallel to $S^1 \times \{0\}$ and there is no holonomy in that case.

Let us denote $\mathcal{F}_{(\tilde{x}_{2s+2}, \dots, \tilde{x}_n)}$ the leaf through the point $(0, 0, \dots, 0, \tilde{x}_{2s+2}, \dots, \tilde{x}_n) \in S^1 \times \mathbb{R}^n$, we get:

$$\mathcal{F}_{(\tilde{x}_{2s+2}, \dots, \tilde{x}_n)} = \{(e^{it_0}, t_1, \dots, t_{2s}, \tilde{x}_{2s+1} = t_{2s+1}, \tilde{x}_{2s+2}, \dots, \tilde{x}_n) \in P^+ / t_0, \dots, t_{2s+1} \in \mathbb{R}\}$$

In the coordinates of the normal form, denote $F_{(x_1, \dots, x_n)}$ the leaf through the point $(0, x_1, \dots, x_n)$, it admits a parametrisation:

$$F_{(x_1, \dots, x_n)} = \{(t_0, (x_i e^{\sum_{j=1 \dots 2s+1} \psi_{i,j} t_j})_{i=1 \dots n}) \in P^+ / t_0 \dots t_{2s+1} \in \mathbb{R}\},$$

where $(\psi_{i,j})$ denotes the inverse matrix of $(\phi_{i,j})$.

REFERENCES

- [Br] O. BRAHIC *Invariants semi-locaux des structures de Poisson*, PhD Thesis, soon available on Arxiv.
- [C1] J.F. CONN, *Normal forms for analytic Poisson structures*, Ann. of Math. (2) **119** (1984) 576-601.
- [C2] J.F. CONN, *Normal forms for smooth Poisson structures*, Ann. of Math. (2) **121** (1985) 565-593.
- [D-Z] J.P. DUFOUR and M. ZHITOMIRSKII, *Classification of non resonant Poisson structure*, J. London Math. Soc. (2) **60** (1999) 935-950.
- [R] O. RADKO, *A classification of generic Poisson structures on a compact oriented surface*, math. SG/0110304.
- [RS] R. ROUSSARIE, *Modèles locaux de champs et de formes*, Asterisque, **30** (1975).
- [W1] A. WEINSTEIN, *The modular automorphism group of a Poisson manifold*, J. Geo. Phys. **23** (1997) 3-4, 379-394.
- [W2] A. WEINSTEIN, *The geometry of Poisson brackets* (Tokyo Universtiy, 1987).